

Antioxidant and antimicrobial activities of Jeok Hasuo (*Polygonum multiflorum* Thunb.) and Baek Hasuo (*Cynanchi wilfordii* Radix) root extracts

Hyunkyung Choi¹, Yuyi Jang², Jun-HyunOh^{1*}

¹Department of Plant and Food Sciences, Sangmyung University, Cheonan 31066, Korea

²SSbio Pharm, Cheonan 31066, Korea

적하수오 및 백하수오 추출물의 항산화 및 항균 활성

최현경¹ · 장유이² · 오준현^{1*}

¹상명대학교 식물식품공학과, ²에스에스바이오팜(주)

Abstract

The objective of this research was to investigate the antioxidant and antimicrobial activities of Jeok Hasuo (*Polygonum multiflorum* Thunb., PM) and Baek Hasuo (*Cynanchi wilfordii* Radix, CW) root extracts. Total phenolic contents of PM and CW root extracts were determined and the antioxidant activities of the root extracts were determined by scavenging activity of diphenylpicrylhydrazyl (DPPH) radicals. The antimicrobial activities against *Staphylococcus aureus* and *Propionibacterium acnes* were determined and expressed as the minimum inhibitory concentration (MIC). The disc diffusion method was also used to determine the zone of inhibition. The butanol extracts of PM and CW roots exhibited greater total phenolic contents (1,212.6 and 1,454.5 mg/g GAE, respectively) than those of ethanol and water extracts. The ethanol (89.0%) and butanol extracts (88.9%) of PM exhibited significantly greater DPPH radical scavenging activities than that of water extracts (73.1%) ($p < 0.05$). Only ethanol extract exhibited an MIC of 0.8 mg/mL against both bacteria. Zones of inhibition started to form when the concentration of extract was greater than 5 mg/disc. The diameters of the zone of inhibition of PM and CW were measured to be 8.9 and 9.2 mm against *S. aureus* and *P. acnes*, respectively, exhibiting the greatest antimicrobial activities among the extracts. This research demonstrated that the PM and CW root extracts possessed not only antioxidant activity but also strong antimicrobial activity against skin-related bacteria.

Key words : antioxidant activity, antimicrobial activity, *Polygonum multiflorum* Thunberg, *Cynanchi wilfordii* Radix

Introduction

Similar traditional herbs have been used in Korea, China, and Japan due to their geographical, cultural, and political, and climatic connections (1). Among these, *Polygonum multiflorum* Thunberg is a popular traditional herbal medicine,

which is called "Hasuo" in Korea ("Heshouwu" in China) (2,3). There are two kinds of "Hasuo", "Jeok Hasuo" and "Baek Hasuo (Back Suo)". Although the names are similar, the origin of the two plants and their botanical characteristics are different. Jeok Hasuo is a root tuber of *Polygonum multiflorum* Thunberg belonging to *Polygonaceae* family. Baek Hasuo is a root tuber of *Cynanchi wilfordii* Radix belonging to *Asclepiadaceae* family (2,3). The difference of botanical characteristics between the two plants lies in the color; Jeok Hasuo is reddish brown, and Baek Hasuo is yellowish brown. In addition, Baek Hasuo contains Baekmilky liquid that is easily differentiated from Jeok Hasuo (4).

*Corresponding author. E-mail : junhyunoh@smu.ac.kr

Phone : 82-41-550-5295, Fax : 82-41-550-5296

Received 25 May 2016; Revised 2 June 2016; Accepted 3 June 2016.

Copyright © The Korean Society of Food Preservation. All rights reserved.

In medicinal studies, Hasuo has been used to treat age-related diseases and for suffering baldness and hair loss (5). In these studies, Hasuo exhibited the various functional properties including blood cholesterol-lowering activity, anti-oxidative activity, anti-inflammatory activity, anti-cancer activity, and neuroprotective effect against ischemia/reperfusion injury (5,6). The major functional components of Baek Hasuo are phosphatidyl choline, phosphatidyl ethanolamine, phosphatidyl inositol, sarcosin, deacylnanchogenin, deacetylmetaplexigenin, hidjoranin, caudatin, penupogenin, methyleugenol, daucosterol, acetovanillone, sucrose, geniposide, and succinic acid (1,7). The major functional components of Jeok Hasuo are emodin, chrysophanol, rhein, physcion, tetrahydroxystilbene, glucopyranoside, monogalloyl ester, resveratrol, polydatin, rhaponticoside, polygonumosides, tricin, rutin, luteolin, quercetin, kaempferol, apigenin, and schizandrin (1,7).

Compared to the extensive research on antioxidant activity of Hasuo, research on antimicrobial activities of Hasuo is limited. Naturally occurring antimicrobial compounds in traditional herbs play an important role in the natural defense or competition systems of living organisms including microorganisms, insects, animals and plants (8). Since Hasuo contains flavones and flavonoid glycosides that may contribute to antimicrobial activity, it can be hypothesized that Hasuo exhibits antimicrobial activity. The pathogen *Staphylococcus aureus* has been a concern not only in food products but also in human skins, because *S. aureus* is one of common pathogens that live on the skin and mucous membranes of humans. In addition, *Propionibacterium acnes* has been known to be one of the major pathogens associated with acnes. Therefore, the main objective of this research was to determine and compare the antioxidant and antimicrobial activities of selected Jeok Hasuo and Baek Hasuo root extracts against skin-related pathogenic bacteria.

Materials and Methods

Materials

Dried roots of Jeok Hasuo (*Polygonum multiflorum* Thunberg, PM) and Baek Hasuo (*Cynanchi wilfordii* Radix, CM) were purchased from a local farm located in Cheonan, South Korea harvested in June 2016. The purchased PM and CW were kept frozen at -20°C prior to extraction. Gallic acid and 1,1-diphenyl-2-picrylhydrazyl (DPPH) were purchased from Sigma Chemical Co. (St. Louis, MO, USA).

Propionibacterium acnes KCTC 3314 and *Staphylococcus aureus* KCTC 3881 were used in this study. *P. acnes* and *S. aureus* were cultured in reinforced clostridial medium (RCM, Difco Laboratories, Detroit, MI, USA) and nutrient broth (Difco Laboratories, Detroit, MI, USA), respectively. *P. acnes* culture was incubated at 37°C for 72 hr under anaerobic conditions in a candle gas jar (GasPak EZ Container Systems, Franklin Lakes, NJ, USA). *S. aureus* culture was incubated in an incubation shaker at 37°C with a 200-rpm shaking for 24 hr. All other reagents and solvents (Sigma Chemical Co., St. Louis, MO, USA) used in this study were analytical grades.

Preparation of Hasuo extracts

The dried roots of PM and CW were cut into small pieces and ground into fine powders using a grinder prior to extraction. Fifty grams of CM or CW were added in 500 mL distilled water, 80% ethanol, and butanol using a soxhlet extractor for 12 hr. After extraction, each extract was filtered through Whatman No. 2 filter paper and the extraction process was repeated twice (9). Then, the solvents of each extract were removed completely using a rotary evaporator (Rikakikal Co., Kyoto, Japan). The extracts were freeze-dried and stored at -20°C prior to use. The extraction yield of the sample was determined using the following equation; Extraction yield (%) = [weight of extract (g)/weight of dried Hasuo (g)] \times 100. Proximate analysis of Hasuo root extracts was conducted following the standard AOAC method (10).

Total phenolic content

The total phenolic contents of Hasuo root extracts were determined following the Folin-Ciocalteu method with minor modifications (11). A portion of 0.50 mL of the diluted extracts was mixed with 0.25 mL Folin-Ciocalteu reagent, followed by the addition of 1.25 mL sodium carbonate (20% aqueous solution). The mixture was kept in the dark for 40 min, and the absorbance was measured at 725 nm using a spectrophotometer (UVmini 1240, Shimadzu, Kyoto, Japan). A standard curve was obtained using gallic acid with selected concentrations of 0.0, 0.02, 0.04, 0.06, 0.08, and 0.1 mg/mL. The total phenolic content of the extracts was expressed as gallic acid equivalents (GAE), which represented the phenolic content as the amount of gallic acid (mg) in 1.0 g sample.

Scavenging activity of diphenylpicrylhydrazyl (DPPH) radical

An aliquot of 1.0 mL of 0.1 mM DPPH radical solution was dissolved in methanol and mixed with 0.5 mL of each

Hasuo extract with a concentration of 0.05 mg/mL or blank methanol for negative control. The reaction solution was mixed, and the absorbance was recorded at 520 nm using a spectrophotometer. Gallic acid (1.5 mg/mL) was used as a standard. The DPPH radical-scavenging activity (%) was calculated by the following equation:

$$\text{Scavenging activity (\%)} = 1 - \left(\frac{\text{Absorbance of sample at 520 nm}}{\text{Absorbance of control at 520 nm}} \right) \times 100$$

The scavenging activity of sample was also expressed as 50% effective concentration (EC₅₀), which represented the concentration of sample exhibiting 50% DPPH radical scavenging activity (11).

Minimum inhibitory concentration of extract against bacteria

The bacterial culture was diluted in the designated broth to contain 8.0 log CFU/mL. The PM and CW root extracts were also diluted with each broth to obtain the concentrations of 0.0, 0.2, 0.4, 0.6, 0.8, and 1.0 mg/mL in the presence of 0.1% Tween 60. The diluted extracts were filtered with sterilized filter (pore size 0.45 μm) before being added to the bacterial culture. An aliquot of 120 μL bacterial culture and 120 μL diluted extracts was incorporated into microplate wells (UV Star, Greiner bio-one, Frickenhausen, Germany). The absorbance was read at 620 nm immediately using a plate reader (Bio-Tek Instrumenting Co., Winooski, VT, USA). The microplate was then incubated according to the cultivation methods described above, and the absorbance was read at 620 nm again after incubation. A minimum inhibition concentration (MIC) at 24 hr was defined as the lowest concentration of antimicrobial that exhibited a complete growth inhibition. The growth inhibition was defined as less than 0.05 absorbance difference (the absorbance of the microplate wells at 24 hr minus the absorbance of the microplate wells at 0 hr) (11).

Disc diffusion method for antimicrobial activity

P. acnes and *S. aureus* were cultivated to possess the concentrations of approximately 10⁶ CFU/mL. One milliliter of each culture was spread on RCM plates for *P. acnes* and on nutrient agar plates for *S. aureus*. Blank paper discs of 6 mm in diameter (BBL, Cockeysville, MD, USA) were impregnated with the selected concentrations of Hasuo root extracts (2, 5, and 10 mg/disc) and placed on the surface of the agar plates. The discs impregnated with Hasuo root extracts were placed under hood to evaporate ethanol and

butanol prior to application on the agar plates. The plate containing *P. acnes* was incubated at 37°C for 72 hr under anaerobic conditions, and the plate containing *S. aureus* was incubated at 37°C for 24 hr. Streptomycin (5 mg/mL) (Dufhefa, Biochemie, Holland) and phosphate-buffered saline (PBS, pH 7.2) were used for positive control and negative control, respectively. After incubation, the diameter of inhibition zone was measured at two cross-sectional points and the average was taken to calculate the area of inhibition zone (12).

Statistical analysis

Each experiment was a completely randomized design. The entire analyses were repeated three times. The treatment factors were selected various solvents (DW, ethanol, and butanol) and two Hasuo root extracts (PM and CW). One way analysis of variance (ANOVA) was performed using SPSS software (version 11.5, SPSS Ins., Chicago, IL, USA). The differences among means were analyzed using Duncan, and the significance level was defined at p<0.05.

Results and Discussion

The proximate analysis of PM and CW roots are presented in Table 1. There was no significant difference in chemical compositions of PM and CW roots except for crude protein contents, however CW root contained a higher amount of crude protein than that of PM. When the Hasuo roots were extracted using different solvents, the water extract exhibited higher extraction yield than those of ethanol extract or butanol extract (Fig. 1). The butanol extracts of PM and CW exhibited the lowest extraction yields among the solvent extracts. However, there was no significant difference in extraction yields between PM and CW in the butanol extraction solvent. Meanwhile, the water extract showed the greatest extraction yield among the solvent extracts.

Table 1. Proximate analysis of *Polygonum multiflorum* Thunberg (PM) and *Cynanchi wilfordii* Radix (CW) roots

	Water (g/100 g)	Crude Protein (g/100 g)	Crude Lipids (g/100 g)	Ash (g/100 g)	Carbohydrates (g/100 g)	Calories (Kcal/100 g)
PM	11.5±0.6	4.25±0.23	0.35±0.17	4.31±0.78	78.91±0.85	326±6
CW	12.2±0.4	7.36±0.51	1.84±0.23	3.45±0.56	72.32±0.45	341±5

Phenolics or polyphenols have received growing attention because of recent findings on their biological activities (13). Most polyphenolic compounds are plant derived secondary




Fig. 1. Extraction yields of *Polygonum multiflorum* and *Cynanchi wilfordii* roots using a Soxhlet extractor.

Different letters (a-c) indicate significant differences at $p < 0.05$.

metabolites and possess antioxidant activities, preventing lipid peroxidation, scavenging free radicals, and chelating redox-active metal ions. Polyphenolic compounds include a wide range of chemicals including flavonoids and proanthocyanidins (14). The total phenolic contents expressed as gallic acid equivalent of PM and CW root extracted with selected solvents are presented in Fig. 2. The phenolic contents of PM and CW were significantly increased in the order of water extract, ethanol extract and butanol extract. The butanol extracts of PM and CW roots contained approximately 1,212.6 and 1,454.5 mg/g GAE, respectively. Wong et al. (15) also investigated the total polyphenolic contents of 60 Chinese medicinal plants, and the authors reported approximately 300 mg/g of total polyphenolic content of PM. The total phenolic contents of butanol extracts from PM and CW exhibited approximately three-fold greater phenolic contents than that




Fig. 2. Total phenolic contents expressed as gallic acid equivalent (GAE) of *Polygonum multiflorum* and *Cynanchi wilfordii* root extracts using selected solvents.

Different letters (a-e) indicate significant differences at $p < 0.05$.

of the water extracts. There was no distinct trend in the total phenolic contents between PM and CW.

The DPPH scavenging activities of PM and CW root extracts with the concentration of 20 mg/mL are presented in Fig. 3. For PM, ethanol extract and butanol extract exhibited 89.0% and 88.8% scavenging activities, respectively, which were significantly higher than that of water extract (73.1%) ($p < 0.05$). For CW, there was no significant difference among the extracts. It was well known that PM possessed strong antioxidant activities *in vitro* and *in vivo* (16-18). However, there is limited information on the functional activities other than antioxidant activities of Hasuo root extracts. In general, chemical compounds responsible for antioxidant activity also possess antimicrobial activities (19). Therefore, it can be anticipated that PM or CW root extracts might exhibit antimicrobial activities.




Fig. 3. DPPH scavenging activities of *Polygonum multiflorum* and *Cynanchi wilfordii* root extracts at the concentration of 20 mg/mL.

Different letters (a-b) indicate significant differences at $p < 0.05$.

The minimum inhibitory concentrations (MICs) of PM and CW root extracts against *P. acnes* and *S. aureus* are presented in Table 2. The water extract and butanol extract of PM and CW did not exhibit any inhibitory effect against the tested bacteria at the tested concentration (1.0 mg/mL). However, the ethanol extracts of PM and CW exhibited MIC of 0.8 mg/mL and 1.0 mg/mL, respectively, against the tested bacteria. The results indicated that PM possessed higher inhibitory effects against tested bacteria than CW against tested bacteria. Although there is limited report on the antimicrobial activity of PM extract, there is no antimicrobial report on CW extract. Hasan et al. (20) investigated the antibacterial activity of *Polygonum hydropiper* root extract, similar to PM, and reported the MIC of 16 μ g/mL against *Staphylococcus aureus*. The authors also indicated that the

antibacterial potency of the plant extract expressed in MIC was stronger against gram-positive bacteria at lower concentration than it was against gram-negative bacteria.

The antimicrobial activities of PM and CW were also determined using a disc diffusion method, and the results are presented in Tables 3 and 4, respectively. Since the preliminary test using the similar ranges for MIC (up to 1.0 mg/disc) did not exhibit any zone of inhibition, the concentrations were increased to 2, 5 and 10 mg/disc. There was no formation of a zone of inhibition at 2 mg/disc-concentration for all of the extracts. However, as the concentrations of extracts were increased, the zones of inhibition were formed, and the diameters of the zones were increased, which was proportional to the concentrations. The ethanol extract possessed the strongest inhibitory effects among the extracts, exhibiting the diameters of inhibition zone ranging from 8.9 to 9.2 mm against tested bacteria. The water extracts exhibited the weakest antimicrobial activities. There was no significant difference between the tested bacteria, presumably because the tested bacteria in this study are gram-positive bacteria.

Table 2. Minimum inhibitory concentration (MIC) of *Polygonum multiflorum* and *Cynanchi wilfordii* root extracts against *P. acnes* and *S. aureus*

Bacteria	Extract	MIC (mg/mL)	
		<i>P. multiflorum</i>	<i>C. wilfordii</i>
<i>Propionibacterium acnes</i>	Water	>1.0	>1.0
	Ethanol	0.8	1.0
	Butanol	>1.0	>1.0
<i>Staphylococcus aureus</i>	Water	>1.0	>1.0
	Ethanol	0.8	1.0
	Butanol	>1.0	>1.0

Kwon (13) also confirmed the antimicrobial activity of PM against both *Staphylococcus epidermidis* and *P. acnes*. The authors could obtain the distinct inhibition zones from the water and 70% ethanol extracts. Hasan et al. (20) reported that chloroform extract of *Polygonum hydropiper* root (150 µg/disc) exhibited a 16-mm zone of inhibition against *S. aureus*. In good agreement with the previous reports, the PM and CW extracts possessed the antimicrobial activities against disease causing bacteria, especially for skin related diseases. *S. aureus* are a concern not only in food products but also in human skins. *S. aureus* are a common type of pathogens that live on human skin and mucous membranes of humans. In addition, *P. acnes* are closely linked to the skin conditions such as acnes. Therefore, the PM or CW root extracts could

be employed to develop skin-associated products and cosmetics due to the antimicrobial activity against skin-related bacteria.

Table 3. The diameter of zone of inhibition of *Polygonum multiflorum* root extracts against *Propionibacterium acnes* and *Staphylococcus aureus*

Bacteria	Extract	Diameter of zone of inhibition (mm)		
		2 mg/disc	5 mg/disc	10 mg/disc
<i>P. acnes</i>	Water	0.0	6.3±0.1	7.5±0.1
	Ethanol	0.0	7.2±0.2	9.1±0.2
	Butanol	0.0	6.4±0.1	7.4±0.1
<i>S. aureus</i>	Water	0.0	6.2±0.1	6.4±0.1
	Ethanol	0.0	7.3±0.1	9.2±0.2
	Butanol	0.0	6.3±0.1	8.4±0.1

Table 4. The diameter of zone of inhibition of *Cynanchi wilfordii* root extracts against *Propionibacterium acnes* and *Staphylococcus aureus*

Bacteria	Extract	Diameter of zone of inhibition (mm)		
		2 mg/disc	5 mg/disc	10 mg/disc
<i>P. acnes</i>	Water	0.0	6.2±0.2	7.2±0.1
	Ethanol	0.0	7.0±0.1	8.9±0.2
	Butanol	0.0	6.3±0.1	7.5±0.1
<i>S. aureus</i>	Water	0.0	6.1±0.1	7.2±0.1
	Ethanol	0.0	7.1±0.2	9.1±0.1
	Butanol	0.0	6.2±0.1	7.9±0.2

요 약

본 논문은 국내 적하수오 및 백하수오 추출물의 항산화능 및 항균활성을 구명하기 위한 목적으로 진행하였다. 각 하수오의 열수, 에탄올, 부탄올 추출물에 대하여, 일반성분을 분석한 후, 항산화능에 대해서는 총 페놀함량 및 DPPH 라디칼 소거능을 측정하였으며, 항균활성에 대해서는 *Staphylococcus aureus* 및 *Propionibacterium acnes*에 대한 최소생육저해농도(minimum inhibitory concentration, MIC) 및 생육저해를 통하여 평가하였다. 적하수오 및 백하수오 부탄올 추출물의 총 페놀함량은 각각 1,212.6 및 1,454.5 mg/g GAE로 다른 유기용매 추출물에 비해 유의적으로 높은 함량을 보였다. 이에 비해 DPPH 라디칼 소거능은 열수 추출물에 비해 에탄올 추출물(89.0%), 부탄올 추출물(88.9%)이 유의적으로 높은 값을 나타내었다. MIC 측정 결과, 적·백하수오 에탄올 추출물만이 0.8 mg/mL로 *S. aureus*와 *P. acnes*에 대해서 항균효과를 나타냈다. 디스크 확산법 측정 결과, 모든 유기용매 추출물이 5 mg/disc 농도부터 생육저해를 형성하였으며, 농도가 증가함에 따라 생육저해환의 크기도 증가하였다. 적하수오 및 백하수오

에탄올 추출물의 조사균주에 대한 생육저해완의 직경은 10 mg/mL 농도에서 각각 8.9 및 9.2 mm를 나타내, 추출물 중 가장 우수한 항균활성을 나타냈다. 본 연구는 하수오 추출물이 항산화 효과 뿐만 아니라 피부 건강과 관련된 세균에 대해 항균활성을 보여줌으로써, 하수오 추출물이 피부 건강을 위한 천연 유래의 기능성 화장품 소재로도 활용될 수 있는 잠재적 가능성을 제시하였다.

Acknowledgement

This work (Grants No. C0276772) was supported by Business for Cooperative R&D between Industry, Academy, and Research Institute funded by Korea Small and Medium Business Administration in 2015.

References

1. Lee BJ, Lee KJ (2015) Discrimination and proper use of *Polygoni multiflori radix*, *Cynanchi wilfordii radix*, and *Cynanchi auriculati radix* in Korea : A descriptive review. Roy Soc Med Int Cong, 2015, 1-7
2. Choi HS, Zhu M, Kim CS, Lee JH (2003) Studies of name and herbal origins of Ha-Soo-Oh. Kor J Orient Med, 9, 81-89
3. Ministry of Food and Drug Safety (2015) The Korean herbal pharmacopoeia XI. Ministry of Food and Drug Safety, Osong, Korea
4. Kim MJ, Kim IJ, Choi SY, Han DH, Kim YH, Lim SC, Kim TJ, Nam SY, Song BH, Oh BU, Park CG (2014) Comparison of *Cynanchum wilfordii*, *C. auriculatum*, *Metaplexis japonica* and *Polygonum multiflorum* by morphological characters. Korean J Medicinal Crop Sci, 22, 113-120
5. Seong ES, Hwang IS, Choi JH, Lee JG, Yoo JH, Ahn YS, Park CB, Yu CY (2015) Enhanced biomass and biological activity of 'Hasuo' (*Polygonum multiflorum* Thunberg) grown under LED light. Aust J Crop Sci, 9, 168-174
6. Na YJ, Lee JH (2014) Physicochemical and antioxidant properties of Yanggaeng with *Cynanchi wilfordii* Radix Powder, J Korean Soc Food Sci Nutr, 43, 1954-1958
7. Seo H, Seo GY, Ko SZ, Park YH (2011) Inhibitory effects of ethanol extracts from *Polygoni multiflori radix* and *Cynanchi wilfordii radix* on melanogenesis in melanoma cells. J Korean Soc Food Sci Nutr, 40, 1086-1091
8. Furukawa T, Eshima A, Kouya M, Takio S, Takano H, Ono K (2002) Coordinate expression of genes involved in catechin biosynthesis in *Polygonum hydropiper* cells. Plant Cell Rep, 21, 385-389
9. Lo KM, Cheung PCK (2005) Antioxidant activity of extracts from the fruiting bodies of *Agrocybe aegerita* var. alba. Food Chem, 89, 533-539
10. AOAC (1996) Official Methods of Analysis. 15th ed, Association of Official Analysis Chemist, Washing DC, p 210-210
11. Mak, YW, Chuah LO, Ahmad R, Bhat R (2013) Antioxidant and antibacterial activities of hibiscus (*Hibiscus rosa-sinensis* L.) and *Cassia* (*Senna bicapsularis* L.) flower extracts. J King Saud Unive-Sci, 25, 275-282
12. Branen J, Davidson PM (2000) Activity of hydrolysed lactoferrin against foodborne pathogenic bacteria in growth media: The effect of EDTA. Lett Appl Microbiol, 30, 233-237
13. Kwon NS (2015) Effectiveness of *Polygonum multiflorum* extracts as whitening, antibacterial, and antiseptic agents. MS Thesis, Daegu Hanny University, Korea, p 21-23
14. Park HJ, Zhang N, Park DK (2011) Topical application of *Polygonum multiflorum* extract induces hair growth of resting hair follicles through upregulating Shh and β -catenin expression in C57BL/6 mice. J Ethnopharmacol, 135, 369-375
15. Wong CC, Li HB, Cheng KW, Chen F (2006) A systematic survey of antioxidant activity of 30 Chinese medicinal plants using the ferric reducing antioxidant power assay. Food Chem, 97, 705-711
16. Chen Y, Wang M, Rosen RT, Ho CT (1999) 2,2-Diphenyl-1-picrylhydrazyl Radical-Scavenging Active Components from *Polygonum multiflorum* Thunb. J Agric Food Chem, 47, 2226-2228
17. Hwang KE (2014) Combined effect of Ganghwa mugwort and ascorbid acid on shelf-life of meat products. MS Thesis, Konkuk University, Korea, p 38-45
18. Lee SV, Choi KH, Choi YW, Hong JW, Baek JU, Choi BT, Shin HK (2014) Hexane extracts of *Polygonum multiflorum* improve tissue and functional outcome following focal cerebral ischemia in mice. Mol Med Rep, 9, 1415-1421
19. Baydar NG, Sagdic O, Ozkan G, Cetin S (2006) Determination of antibacterial effects and total phenolic contents of grape (*Vitis vinifera* L.) seed extracts. Int J Food Sci Technol, 41,799-804
20. Hasan MF, Das R, Khan A, Mossain MS, Rahman M (2009) The determination of antibacterial and antifungal activities of *Polygonum hydropiper* (L.) root extract. Adv Biologic Res, 3, 53-56